



Dr Veena Aggarwal
Consultant, Womens' Health
CMD and Group Editor-in-Chief,
IJCP Group & Medtalks
Trustee, Dr KK's Heart Care
Foundation of India

Risk Factors for Postmenopause Depressive Symptoms

Women with depressive symptoms preceding the final menstrual period (FMP) are at greater risk of developing depression during the postmenopause, according to results from the Study of Women's Health Across the Nation (SWAN) recently published in the journal *Menopause*.¹

A total of 1,551 middle-aged participants from the Study of Women's Health Across the Nation were included in the present study. A longitudinal analysis of the data on depressive symptoms gathered from 1996 to 2017 among the selected women was longitudinally analyzed over 19 years (median) to examine depressive symptoms during postmenopause and the contribution of depressive symptom trajectories before the FMP. The study also examined the impact of psychosocial/health factors on the postmenopause depressive symptoms.

Compared to premenopause stage, the probability of depressive symptom scores of 16 or higher on the Center for Epidemiologic Studies Depression Scale (CES-D) was greater during postmenopause with odds ratio (OR) of 1.49, but not during the menopausal transition or perimenopause. Based on the scores, three pre-FMP groups were defined: Group 1 (47.7%), consistently low scores, Group 2 (39.9%), moderate scores below the high depressive symptom threshold and Group 3 (12.4%), consistently high scores.¹ Women with moderate scores on the depression scale were nearly threefold more

likely to have higher scores when postmenopausal with OR of 2.62, whereas women who scored high on the depression scale were at nearly sevenfold greater risk of higher scores during the postmenopause with OR of 6.88.

History of severe anxiety, sleep problems, vasomotor symptoms, low social support and childhood trauma/maltreatment before reaching the FMP were also identified as contributing to higher odds of depressive symptoms during postmenopause.

World Menopause Day is held every year on 18th October and the theme for this year is "Cognition and Mood". Hot flashes and night sweats are typically associated with menopause. However, the symptoms of menopause are wide-ranging. They include mood problems among several others. Earlier studies have shown that mood problems may adversely affect cognition. Hence, women approaching menopause should be regularly screened for risk factors predisposing to depressive symptoms later on and provided with appropriate treatment for a better quality of life.

REFERENCE

1. Kravitz HM, Colvin AB, Avis NE, Joffe H, Chen Y, Bromberger JT. Risk of high depressive symptoms after the final menstrual period: the Study of Women's Health Across the Nation (SWAN). *Menopause*. 2022;29(7):805-15.

